

# Dosage Chart

## Acetaminophen (Tylenol® or another brand)

Give every 6 hours as needed.(Also available in suppositories; use the same number of mg.)

**\*Do not give more than 4 doses in 24 hours.**

Weight in pounds (lbs.)	Elixir = 160 mg/5ml	Chewable 1 tablet = 80 mg	Jr. strength 1 caplet = 160 mg	Reg. Strength 1 tablet = 325mg
6-11 lbs.	1.25 ml	-----	-----	-----
12-17 lbs.	2.5 ml	-----	-----	-----
18-23 lbs.	3.75 ml	-----	-----	-----
24-35 lbs.	5 ml	2 tablets	-----	-----
36-47 lbs.	7.5 ml	3 tablets	-----	-----
48-59 lbs.	10 ml	4 tablets	2 caplets	1 tablet
60-71 lbs.	12.5 ml	5 tablets	2½ caplets	1 tablet
72-95 lbs.	15 ml	6 tablets	3 caplets	1½ tablets
96+ lbs.	-----	-----	4 caplets	2 tablets

## Ibuprofen (Advil® Motrin® or another brand)

Give every 6 to 8 hours as needed; always with food.

Other strengths available by prescription.

**\*Do not give more than 4 doses in 24 hours.**

Weight in pounds (lbs)	Infant drops (40mg/1 ml) Infants > 6 month only!	Children's liquid 100 mg/5ml	Chewable Tablets 1 tablet = 100 mg	Tablets 1 tablet = 200 mg
11-21 lbs.	1.25ml.	2.5 ml	½ tablet	-----
22-32 lbs.	2.5ml	5 ml	1 tablet	-----
33-43 lbs.	-----	7.5 ml	1 ½ tablets	-----
44-54 lbs.	-----	10 ml	2 tablets	1 tablet
55-65 lbs.	-----	12.5 ml	2 ½ tablets	-----
66-87 lbs.	-----	15 ml	3 tablets	1½ tablets
88+ lbs.	-----	20 ml	4 tablets	2 tablets

## Questions?

This sheet is not specific to your child but provides general information. If you have any questions, please call your clinic or pharmacy.